

## CLAIMS

1. A method comprising:

a) providing:

i) a first recombinant vector, comprising in operable combination:

- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest;
- 3) adenovirus packaging sequence linked to one of said inverted terminal repeats; and
- 4) an adeno-associated virus terminal repeat sequence operably linked to said 3' end of said nucleotide sequence of interest,

wherein said first vector lacks a second adeno-associated virus terminal repeat sequence, and lacks one or more adenovirus early gene region selected from E1, E2, E3, and E4 gene region; and

ii) a cell capable of expressing said one or more adenovirus early gene which is lacking from said first vector;

b) introducing said first vector into said cell to produce a transformed cell; and

c) culturing said transformed cell under conditions such that a second vector is produced, said second vector selected from:

- i) a third vector, comprising in operable combination:
  - 1) adeno-associated virus terminal repeat DD sequence;

2) first and second inverted copies of a nucleotide sequence of interest flanking said adeno-associated virus terminal repeat-DD sequence;

3) left and right inverted terminal repeats of adenovirus flanking said first and second inverted copies of said nucleotide sequence of interest; and

4) an adenovirus packaging sequence linked to one of said inverted terminal repeats, and

ii) a fourth vector, comprising in operable combination:

1) a nucleotide sequence of interest having a 5' end and a 3' end;

2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest; and

3) an adenovirus packaging sequence linked to one of said inverted terminal repeats.

2. The method of Claim 1, wherein said cell is capable of expressing one or more Rep proteins, and said culturing results in expression of said one or more Rep proteins.

3. The method of Claim 1, wherein said second vector is encapsidated.

4. The method of Claim 3, further comprising d) recovering said encapsidated second vector.

5. The method of Claim 4, further comprising e) purifying said recovered encapsidated second vector.

6. The method of Claim 5, further comprising e) administering said purified encapsidated second vector to a host cell.

7. The method of Claim 6, wherein said administering is under conditions such that said nucleotide sequence of interest in said encapsidated second vector is expressed.

8. The method of Claim 6, wherein said host cell is a cultured cell.

9. The method of Claim 6, wherein said host cell is comprised in a mammal.

10. The method of Claim 9, wherein said mammal is selected from mouse and human.

11. The method of Claim 2, wherein expression of one or more Rep proteins is inducible.

12. A method comprising:

a) providing:

i) a first recombinant vector, comprising in operable combination:

1) a nucleotide sequence of interest having a 5' end and a 3' end;

2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest;

3) adenovirus packaging sequence linked to one of said inverted terminal repeats; and

- 4) an adeno-associated virus terminal repeat sequence operably linked to said 3' end of said nucleotide sequence of interest,

wherein said first vector lacks a second adeno-associated virus terminal repeat sequence, and lacks one or more adenovirus early gene region selected from E1, E2, and E4 gene region;

- ii) a cell capable of expressing one or more Rep proteins; and

- iii) helper adenovirus;

b) introducing said first vector and genome of said helper adenovirus into said cell to produce a transformed cell; and

c) culturing said transformed cell under conditions such that said transformed cell expresses said one or more Rep proteins, and a second vector is produced, said second vector selected from:

- i) a third vector, comprising in operable combination:

- 1) adeno-associated virus terminal repeat DD sequence;

- 2) first and second inverted copies of a nucleotide sequence of interest flanking said adeno-associated virus terminal repeat-DD sequence;

- 3) left and right inverted terminal repeats of adenovirus flanking said first and second inverted copies of said nucleotide sequence of interest; and

- 4) an adenovirus packaging sequence linked to one of said inverted terminal repeats, and

- ii) a fourth vector, comprising in operable combination:

- 1) a nucleotide sequence of interest having a 5' end and a 3' end;

- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest; and
- 3) an adenovirus packaging sequence linked to one of said inverted terminal repeats

13. The method of Claim 12, wherein said cell lacks expression of said one or more adenovirus early gene region which is lacking from said first vector.

14. A method comprising:

a) providing:

i) a first recombinant vector, comprising in operable combination:

- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest;
- 3) adenovirus packaging sequence linked to one of said inverted terminal repeats; and
- 4) an adeno-associated virus terminal repeat sequence operably linked to said 3' end of said nucleotide sequence of interest,

wherein said first vector lacks a second adeno-associated virus terminal repeat sequence, and lacks one or more adenovirus early gene region selected from E1, E2, and E4 gene region;

- ii) a cell capable of expressing said one or more adenovirus early gene which is lacking from said first vector; and
- iii) adeno-associated virus;

- b) introducing said first vector and genome of said adeno-associated virus into said cell to produce a transformed cell; and
- c) culturing said transformed cell under conditions such that a second vector is produced, said second vector selected from:

- i) a third vector, comprising in operable combination:
- 1) adeno-associated virus terminal repeat DD sequence;
  - 2) first and second inverted copies of a nucleotide sequence of interest flanking said adeno-associated virus terminal repeat-DD sequence;
  - 3) left and right inverted terminal repeats of adenovirus flanking said first and second inverted copies of said nucleotide sequence of interest; and
  - 4) an adenovirus packaging sequence linked to one of said inverted terminal repeats, and
- ii) a fourth vector, comprising in operable combination:
- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
  - 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest; and
  - 3) an adenovirus packaging sequence linked to one of said inverted terminal repeats.

15. A method comprising:

- a) providing:
- i) a first recombinant vector, comprising in operable combination:

- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest;
- 3) adenovirus packaging sequence linked to one of said inverted terminal repeats; and
- 4) an adeno-associated virus terminal repeat sequence operably linked to said 3' end of said nucleotide sequence of interest,

wherein said first vector lacks a second adeno-associated virus terminal repeat sequence, and lacks adenovirus E3 early gene region; and

- ii) a cell;
- b) introducing said first vector into said cell to produce a transformed cell; and
- c) culturing said transformed cell under conditions such that a second vector is produced, said second vector selected from:
  - i) a third vector, comprising in operable combination:
    - 1) adeno-associated virus terminal repeat DD sequence;
    - 2) first and second inverted copies of a nucleotide sequence of interest flanking said adeno-associated virus terminal repeat-DD sequence;
    - 3) left and right inverted terminal repeats of adenovirus flanking said first and second inverted copies of said nucleotide sequence of interest; and
    - 4) an adenovirus packaging sequence linked to one of said inverted terminal repeats, and
  - ii) a fourth vector, comprising in operable combination:

- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest; and
- 3) an adenovirus packaging sequence linked to one of said inverted terminal repeats.

16. The method of Claim 15, wherein said cell is capable of expressing one or more Rep proteins, and said culturing results in expression of said one or more Rep proteins.

17. A method comprising:

- a) providing:
  - i) a first recombinant vector, comprising in operable combination:
    - 1) a nucleotide sequence of interest having a 5' end and a 3' end;
    - 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest;
    - 3) adenovirus packaging sequence linked to one of said inverted terminal repeats; and
    - 4) an adeno-associated virus terminal repeat sequence operably linked to said 3' end of said nucleotide sequence of interest,

wherein said first vector lacks a second adeno-associated virus terminal repeat sequence, and wherein said nucleotide sequence of interest in said first vector comprises adeno-associated virus rep gene region; and



- 5
- ii) a cell;
- b) introducing said first vector into said cell to produce a transformed cell; and
- c) culturing said transformed cell under conditions such that said transformed cell expresses one or more Rep proteins, and a second vector is produced, said second vector selected from:
- 10
- i) a third vector, comprising in operable combination:
- 1) adeno-associated virus terminal repeat DD sequence;
- 2) first and second inverted copies of a nucleotide sequence of interest flanking said adeno-associated virus terminal repeat-DD sequence;
- 15
- 3) left and right inverted terminal repeats of adenovirus flanking said first and second inverted copies of said nucleotide sequence of interest; and
- 4) an adenovirus packaging sequence linked to one of said inverted terminal repeats, and
- ii) a fourth vector, comprising in operable combination:
- 20
- 1) a nucleotide sequence of interest having a 5' end and a 3' end;
- 2) left and right inverted terminal repeats of adenovirus flanking said nucleotide sequence of interest; and
- 25
- 3) an adenovirus packaging sequence linked to one of said inverted terminal repeats.

18. The method of Claim 17, wherein said first vector lacks one or more adenovirus early gene region selected from E1, E2, and E4 gene region, and said cell

is capable of expressing said adenovirus early gene region which is lacking from said first vector.

19. The method of Claim 17, wherein said first vector lacks adenovirus E3 gene region.

5

1000 2000 3000 4000 5000 6000 7000 8000 9000 10000 11000 12000 13000 14000 15000 16000 17000 18000 19000 20000 21000 22000 23000 24000 25000 26000 27000 28000 29000 30000 31000 32000 33000 34000 35000 36000 37000 38000 39000 40000 41000 42000 43000 44000 45000 46000 47000 48000 49000 50000 51000 52000 53000 54000 55000 56000 57000 58000 59000 60000 61000 62000 63000 64000 65000 66000 67000 68000 69000 70000 71000 72000 73000 74000 75000 76000 77000 78000 79000 80000 81000 82000 83000 84000 85000 86000 87000 88000 89000 90000 91000 92000 93000 94000 95000 96000 97000 98000 99000 100000